

# Antarctic tardigrades a biological model for geobiology and astrobiology studies: a review

<sup>1</sup>Rosa Acevedo-Barrios<sup>1</sup>; Wendy Miranda-Castro<sup>1</sup>; Diana González<sup>1</sup>; Jorgelina Pasqualino<sup>2</sup>; Irina P. Tirado-Ballestas<sup>3</sup>; María Angélica Leal<sup>4</sup>, David Tovar<sup>4</sup>

<sup>1</sup> Grupo de Estudios Químicos y Biológicos, Dirección de Ciencias Básicas, Universidad Tecnológica de Bolívar, POB 130001, Cartagena de Indias D. T. y C., Colombia, [racevedo@utb.edu.com](mailto:racevedo@utb.edu.com), [wmiranda@utb.edu.co](mailto:wmiranda@utb.edu.co), [gonzalezd@utb.edu.co](mailto:gonzalezd@utb.edu.co), [drubiano@utb.edu.co](mailto:drubiano@utb.edu.co)

<sup>2</sup> Grupo de Investigación en Sistemas Ambientales e Hidráulicos (GISAH), Escuela de Ingeniería Arquitectura & Diseño, Universidad Tecnológica de Bolívar, Cartagena, Colombia, [cdiaz@utb.edu.co](mailto:cdiaz@utb.edu.co); [jpasqualino@utb.edu.co](mailto:jpasqualino@utb.edu.co)

<sup>3</sup> GENOMA Group, Health Sciences Department, Universidad del Sinú, Santillana Campus, Cartagena, Colombia, [irina.tirado@unisinu.edu.co](mailto:irina.tirado@unisinu.edu.co)

<sup>4</sup> Planetary Sciences and Astrobiology Research Group (GCPA), Universidad Nacional de Colombia, Bogotá, Colombia, [maleall@unal.edu.co](mailto:maleall@unal.edu.co), [dftovarr@unal.edu.co](mailto:dftovarr@unal.edu.co)

**Abstract**– *Tardigrades (commonly known as "water bears") are microscopic animals whose bodies are usually less than 1 mm long, they are microscopic bilaterian organisms that belong to the phylum Tardigrada. These organisms are best known for initiating and maintaining a state of dormancy known as cryptobiosis. This ability allows them to survive in unfavorable environments and to inhabit places characterized by extreme temperatures, variable water availability, etc. (e.g., Antarctica). Antarctica is of great scientific interest, as the extreme environmental conditions require unique adaptive traits expressed by the organisms inhabiting this region. The presence of tardigrades in Antarctica has been scientifically documented, and their characteristics could be strategic in developing geobiological, astrobiological, and other areas of knowledge.*

**Keywords**– *biodiversity, cryptobiosis, evolutionary adaptations, extreme environment, polar ecology.*

## I. INTRODUCTION

Limno-terrestrial tardigrades are microscopic bilaterian organisms that belong to the phylum Tardigrada within the larger protostome superclade called Ecdysozoa. [1]. They are also known as water bears or moss piglets [2]. These organisms are close relatives of arthropods and Onychophora [3], with body sizes from 50  $\mu\text{m}$  to 1200  $\mu\text{m}$ , and eight legs [4], [5]. The body of a tardigrade is composed of five segments, which include a head and four trunk segments. Typically, they consist of only about 1,000 cells [6].

Over the years, the number of identified tardigrade species has grown significantly since the initial species description by Schultz [7]. Currently, there are more than 1400 acknowledged species within the phylum [8].

The semi-terrestrial tardigrades are found in soil, sediments, bryophytes, lichens, algae (where they are active when a water film surrounds the substrate), and lotic and lentic environments. They can be found from low to high altitudes and latitude, from tropical, temperate, desert, and polar habitats [9]. Additionally, tardigrades having an estimated life

span of three to 30 months, without including their latency or cryptobiosis period [9].

The majority of tardigrades rely on water for growth and reproduction, certain species found in limno-terrestrial habitats possess the remarkable capability to endure drought [10] and tolerate a variety of extreme environmental conditions at any stage of their life because of their ability to enter cryptobiosis, a process in which metabolic activity is temporarily suppressed [11], [12], [13]. This trait enables water bears to survive in extreme environments such as Antarctica [14], [15].

The extreme climatic conditions of Antarctica are of high scientific interest, and has contribute to organisms developing or displaying unique evolutionary adaptations to the environmental extremes that characterise Antarctica: low temperatures, desiccation, high salinity (in coastal regions), high solar irradiance [16]. Research on Antarctic tardigrades started in the early 20th century, but was sporadic and associated with expeditions (e.g. Richters 1904a; Murray 1907; Richters 1908; Murray 1910 [17], [18], [19], [20]). The knowledge of Antarctic tardigrades is still relatively poor. In 2005, Convey and McInnes reported on the presence of tardigrade species on ten of the 11 major islands of the Antarctic Peninsula, and including previous reports, approximately 55 species have been documented in the Antarctic region.

The Antarctic Peninsula has a higher variety of tardigrade species than continental Antarctica [21]. However, knowledge of the biogeographic distribution, abundance, and diversity of tardigrades in Antarctica remains limited [22], making them an attractive research topic. [23], [24]. The relationships between tardigrades, geology, and space research have been even less studied, but with a great perspective of development as a research topic. For this reason, the present article aims to review the research potential of limno-terrestrial tardigrades in Antarctica as a model for geobiological and astrobiological studies.

## II. MATERIALS AND METHODS

To analyse the geographic distribution of tardigrade species reported in Antarctica, academic texts were collected and reviewed from different databases from 2022 to 2024. This literature review was conducted using articles found in ResearchGate, SpringerLink, Csiro Publishing, Scopus and Google scholar. The search was conducted using “Tardigrade”, “Antarctic” and “Antarctica” as keywords. Also, articles found in references section of the articles selected were manually searched and included in the review. Peer reviewed articles presenting information about the occurrence of limno-terrestrial tardigrades in Antarctic were added, including their isolation and identification process.

## III. TARDIGRADES AND CRYPTOBIOSIS

Tardigrades can be categorized into two primary evolutionary lineages: eutardigrades and heterotardigrades. Among these, the heterotardigrades exhibit greater diversity in terms of species and characteristics [25].

Tardigrades do not possess specialized respiratory organs; instead, gas exchange occurs through diffusion across the epidermis and cuticle [5]. Similarly, they lack circulatory organs like a heart, and instead rely on a fluid-filled body cavity for circulatory functions. This cavity accommodates storage cells of varying sizes and quantities, which move passively along with the animal's movements [26]. The storage cells play important roles in nutritional maintenance, vitellogenesis, and potentially contribute to immune defense mechanisms [27]. Tardigrades possess longitudinal muscles and excretory systems. Males, females, and hermaphrodites capable of self-fertilization exhibit a singular dorsal sack-like gonad. In certain species, reproduction solely occurs through parthenogenesis, with only female individuals present [28].

Tardigrades have the capacity to undergo cryptobiosis, a state of reversible standstill often referred to as latent or hidden life, is prevalent across various kingdoms of life [29]. Among metazoans, nematodes, rotifers, and tardigrades possess the remarkable ability to enter cryptobiosis at any stage of their life cycle, including as eggs, juveniles, and adults [30].

The entry into the cryptobiotic state involves a sequence of anatomical and physiological transformations. The presence of a water film is crucial for tardigrades to maintain their physiologically active state [31]. When faced with desiccation, during which tardigrades can lose over 95% of their water content, they undergo longitudinal contraction. In this process, they retract their head and legs, assuming a dormant and barrel-shaped structure known as a “tun” [32]. In this state, oxygen consumption almost stops, and the metabolic rate decreases considerably [33], [34], [35]. Nonetheless, there is compelling evidence suggesting that certain species of tardigrades can endure substantial levels of environmental stress even while in an active state [36]. Cryptobiosis can be triggered by various extreme conditions, leading to different

stages of cryptobiotic dormancy. Notably, stressors such as; water loss (desiccation), increased external osmotic pressure (osmobiosis), freezing (cryobiosis), lack of oxygen (anoxibiosis), and exposure to environmental toxins (chemobiosis), can induce a state of quiescence [37], [38], [39], [40], [41], [42], [43]. Anhydrobiosis, osmobiosis, and potentially chemobiosis are cryptobiotic states characterized by the formation of tuns. The ability to form tuns is observed in some existing tardigrade lineages, indicating that it is an ancient and shared characteristic [44].

The interest in the investigation of tardigrade physiology is growing because of their ability to withstand a diverse range of abiotic challenges. These challenges encompass desiccation [31], [38], elevated levels of ionizing and UV radiation [45], [46], the vacuum of space [47], [48], contact with harmful heavy metals and metalloids [49], [50], extreme temperatures spanning from very low to high [40], [51], [52], and varying atmospheric conditions, including both low and high pressures [41], [53], and even oxygen deprivation [54], [55].

Despite the limited understanding of the cryptobiosis process, the mounting evidence indicates that tardigrades produce a diverse array of molecules with bioprotective properties [56], [57]. These molecules include proteins, such as DNA repair systems, as well as carbohydrates, among other substances [58]. For instance, there is a unique protein called Damage suppressor (Dsup) found in tardigrades that plays a crucial role in safeguarding DNA. This tardigrade-specific protein can bind to DNA and nucleosomes [59], [60], [61].

When introduced into human cultured cells, Dsup effectively mitigates the occurrence of DNA breaks induced by radiation and reactive oxygen species (ROS). As a result, the expression of Dsup enhances the survival of cells exposed to a semi-lethal dose of X-ray irradiation [62]. Tardigrades possess additional unique proteins called Cytosolic Abundant Heat-Soluble (CAHS) and Secretory Abundant Heat-Soluble (SAHS) proteins. These proteins have been associated with the tardigrades' capacity to endure desiccation, enabling their survival in extremely dry conditions [63]. These biological capabilities and the ability to enter a cryptobiosis state facilitate and increase the survival and presence of tardigrades in Antarctica.

## IV. ANTARCTIC TARDIGRADES

According to the literature, the presence of tardigrades in Antarctica was first validated in 20<sup>th</sup> century [64]. Richters [65], was the first to report the presence of tardigrades in Antarctica. Until 1962, only four genera had been identified, namely, *Echiniscus*, *Pseudechiniscus*, *Hypsibius*, and *Milnesium*. These genera were mostly collected from the South Orkney Islands, Victoria Land, South Shetland, and other locations in Antarctica [66]. In Larsemann Hills, East Antarctica, surveyed in 1987, five genera and six species of Tardigrada were identified, and dynamics linked to dispersal capabilities of tardigrades in Antarctica were proposed [67].

In 1988, A total of 28 species of tardigrades were known from the Vestfold Hills in Antarctica (Miller et al., 1988). These species have been reported in Gaussberg [69], Syowa, Queen Maud Land (Japanese base) [70], Vestfold Hills (Davis Base), and the Clark Peninsula (Wilkes Base) [67], [71]. Understandably, the first reports of tardigrades were of locations near bases; therefore, most knowledge of tardigrades in Antarctica was focused on East Antarctica and the Antarctic Peninsula [67], [72], [73]. There have also been reports of tardigrades in Enderby Land, Coast of Alasheev Bight, Antarctic Peninsula, and South Shetland Islands - King George Island.

At the end of the 20th century, Dastych redescribed the species *Hypsibius antarcticus* to prevent it being confused with *Hypsibius arcticus* and reassigned all Antarctic tardigrades cited as *Hypsibius arcticus* in the literature to *Hypsibius antarcticus* [74], [75]. Pilato & Binda [76] revised the genus to *Acutuncus antarcticus*. Other species of tardigrades reported during the 20th century have been redescribed due to the improving taxonomic methods in the last decades and the increasing interest in studying these organisms to help classify them according to their similarities. For example, the species *Echiniscus pseudowendti* was reassigned to *Claxtonia pseudowendti*. In summary, around 35 tardigrade species were documented in the 20th century such as *Pseudechiniscus suillus*, *Acutuncus antarcticus*, *Ramajendas frigidus*, *Diphascon chilensis*, *Dip. pingue*, *Dip. polare*, *Dip. dastychi*, and *Dip. victoriae* [77].

In 2000, a new species of eutardigrade, *Diphascon (Adropion) tricuspdatum* (now *Adropion tricuspdatum*) was found with *Acutuncus antarcticus* in a small lake in the Crater Cirque (Victoria Land) [78]. The following year, more information on tardigrades obtained from the 1977 to 78 Australian Museum Expedition was published, expanding on information highlighting the evolutionary relationships of tardigrades in Antarctica and confirming the biogeographic distinctiveness of *Adropion tricuspdatum*.

TABLE I  
TARDIGRADES DOCUMENTED IN ANTARCTICA DURING THE 20<sup>TH</sup>  
CENTURY

	Genera	Species	Reference
1	<i>Acutuncus</i>	<i>antarcticus</i>	[18], [69], [72], [74], [80], [81], [82]
2	<i>Adropion</i>	<i>greveni</i>	[72]
3	<i>Barbaria</i>	<i>jenningsi</i>	[72], [80], [83]
4	<i>Claxtonia</i>	<i>pseudowendti</i>	[74]
5	<i>Dastychius</i>	<i>improvisus</i>	[74]
6	<i>Dianeia</i>	<i>papillifer</i>	[84], [85]
7	<i>Diphascon</i>	<i>alpinum</i>	[72], [74], [86]
8	<i>Diphascon</i>	<i>chilensis</i>	[83], [87]
9	<i>Diphascon</i>	<i>mirabile</i>	[74]
10	<i>Diphascon</i>	<i>pingue</i>	[72], [81]
11	<i>Diphascon</i>	<i>puniceum</i>	[86]
12	<i>Diphascon</i>	<i>sanae</i>	[80], [88], [81]

13	<i>Diphascon</i>	<i>polare</i>	[77]
14	<i>Diphascon</i>	<i>dastychi</i>	[77]
15	<i>Diphascon</i>	<i>victoriae</i>	[77]
16	<i>Echiniscus</i>	<i>kergeuelensis</i>	[84]
17	<i>Echiniscus</i>	<i>darienae</i>	[80]
18	<i>Echiniscus</i>	<i>punctus</i>	[85]
19	<i>Grevenius</i>	<i>asper</i>	[85]
20	<i>Grevenius</i>	<i>laevis</i>	[85]
21	<i>Hebesuncus</i>	<i>schusteri</i>	[74]
22	<i>Hebesuncus</i>	<i>conjungens</i>	[82]
23	<i>Hypsibius</i>	<i>convergens</i>	[89]
24	<i>Hypsibius</i>	<i>dujardini</i>	[72]
25	<i>Hypsibius</i>	<i>simoizumii</i>	[90]
26	<i>Mesobiotus</i>	<i>blocki</i>	[80]
27	<i>Mesobiotus</i>	<i>fuciger</i>	[72], [91]
28	<i>Mesobiotus</i>	<i>harmsworthi</i>	[84], [87]
29	<i>Milnesium</i>	<i>tardigradum</i>	[72], [85], [82]
30	<i>Minibiotus</i>	<i>weinerorum</i>	[67], [88]
31	<i>Minibiotus</i>	<i>stuckenbergi</i>	[92], [88]
32	<i>Mixibius</i>	<i>saracenus</i>	[82]
33	<i>Oreella</i>	<i>mollis</i>	[85]
34	<i>Pseudechiniscus(Meridioniscus )</i>	<i>novaezeelandiae</i>	[68], [93]
35	<i>Pseudechiniscus(Pseudechiniscus)</i>	<i>suillus</i>	[85], [93]
36	<i>Ramajendas</i>	<i>heatwolei</i> sp.	[80]
37	<i>Ramajendas</i>	<i>frigidus</i>	[94]
38	<i>Ramajendas</i>	<i>renaudi</i>	[72], [95]
39	<i>Ramazottius</i>	<i>oberhaeuseri</i>	[96], [97]

Previous reports and descriptions of tardigrades were usually studies focusing on which tardigrades are to be found in Antarctica; however, more studies on abundance, frequency, species richness, and their correlation with others species have been published. A study conducted in the nunataks of the Schirmacher Oasis found that the highest densities of tardigrades were observed at sites with mosses, lichens, liverworts, or algae [99]. The most abundant and frequently occurring tardigrade genera were found to be *Mesobiotus* and *Hebesuncus* [99]. In addition, studies on tardigrade colonization and dynamics have also been published (e.g. Smykla et al. 2012). Results showing post-glacial and Holocene dynamics were obtained through the assessment of tardigrade eggs and exuviae from Antarctic lake sediments and paleosediments [101].

A new species of Heterotardigrada, *Echiniscus corrugicaudatus* (now *Claxtonia corrugicaudatus*) was discovered by McInnes (2010) in the nunataks of Ellsworth Land, West Antarctica. In 2012, *Milnesium antarcticum* was documented in inland Antarctica for the first time, more specifically in Victoria Land [100], [103]. The *Mil. antarcticum* specie were also found with *Acutuncus antarcticus* and were present in 23 of 41 samples. The occurrence of these two tardigrades species in Victoria Land is linked to the high water availability in the soil [100];

overall, abiotic conditions influence the abundance of these animals [104].

The location near Syowa station is one of places in continental Antarctica [105], [106]; therefore, an update of tardigrade diversity was reported in this area, and three new species were documented: *Claxtonia pseudowendti*, *Hebesuncus ryani*, and *Pseudechiniscus* sp.§ [106]. Another location is Victoria Land, where numerous species of tardigrades have been reported. Pilato et al. (2017) included new information on the Tardigrada biodiversity, updating the presence of *Diphascion sanae* and reporting two new species, namely, *Mixibius felix* and *Milnesium validum*.

In recent years, molecular characterisation in conjunction with morphological descriptions, i.e. integrative taxonomy, has been used to facilitate the identification of new taxa. The species *Mopsechiniscus franciscaae* [108], *Cryoconicus antiarktos*, and *Ramazottius sabatiniae* [109] were identified using this technique.

With the advent of technological advances, modern molecular studies have been applied to understanding the evolutionary relationships of Antarctic tardigrades. For example, an 18S rRNA sequencing was performed to confirm a culture of tardigrades from near Syowa station, and the presence of *Acutuncus antarcticus* was confirmed. Older records had confused this species with *Hypsibius articus* [105]. The use of molecular techniques to identify mitochondrial operational taxonomic units (OTUs), resulted in potentially more putative species in the genera *Acutuncus*, *Milnesium*, and *Echiniscus* than have been reported using older morphological methods [110].

These new tools used for identification methods allow a faster classification [111], [112]. For example, various species have been redescribed, such as *Hypsibius dujardini* [113], *Hypsibius murray* [114], and the genus *Ramajendas* [115]. However, taxonomic studies of Antarctic tardigrades have developed slowly and the diversity and distribution of these animals in this extreme continent remains largely unknown. This is mainly due to difficulties associated with working in the extreme climatic conditions of Antarctica, distance, transport and mobility, which make most of the continent inaccessible and prevent comprehensive studies (Fig 1). A list of tardigrade species documented since 2000 is provided in.




Fig. 1 Location of tardigrades collected from different locations of the Antarctic continent.

## V. GEOBIOLOGICAL RELATIONSHIPS

Geological processes can be understood from the established biological relationships; in turn, geological dynamics can model diverse biological processes. This relationship between life and a planet's geological history is known as geobiology [116]. Depending on the biological scale at which one works, the geological processes that may impact will vary [117]. Biogeochemical cycles are one of the most important relationships between microorganisms and geology [117], [118].

In addition, changes such as sea level, volcanic activity, or mineralogical composition of rocks directly affect the distribution and diversity of microorganisms [119], [120]. Some of the most determinant geological parameters for the presence and abundance of certain species of microorganisms are soil type, water availability, and the presence of certain minerals [121], [122], [123]. On the other hand, some microorganisms can serve as bioindicators of geological changes [124].

For most microorganisms, it has been evidenced that they tend to be more abundant and diverse in soils rich in organic matter [117], [125]. However, in the case of tardigrades studied in Antarctic sectors, it has been found that the determining factor was the water content in the soil, this taking into account that in Antarctic sectors with ornithogenic soils, major presence and diversity of tardigrades was observed [100], [126].

On the other hand, it has been identified that limnoterrestrial tardigrades could be an indicator of geological history. Tardigrades though not part of the fossil record and, until recently, thought to be ubiquitous due to their aerial dispersal and cryptobiosis [127], have shown that they are associated with specific geological and environmental characteristics and with the scale of geological time (Pilato and Binda 2001; McInnes and Philip 2007; Guidetti et al. 2017). Finally, tardigrades could be diverse in mineralogical environments, being organisms that can tolerate toxicity, contribute to nutrient cycling, and use nutrient diversity [130], [131].

#### VI. ASTROBIOLOGICAL PERSPECTIVES

Considering the characteristics mentioned above, which allow tardigrades to survive in extreme conditions, they have been considered a possible model organism for the search for life outside Earth [47]. Tests on several tardigrade species suggest they could survive in conditions similar to those found on other planets and moons [132].

TABLE II  
ANTARCTIC TARDIGRADES DISCOVERED IN THE 21<sup>ST</sup> CENTURY

	Genera	Species	Reference
1	<i>Adropion</i>	<i>tricuspidatum</i>	[78]
2	<i>Bryodelphax</i>	<i>olszanowskii</i>	[133]
3	<i>Claxtonia</i>	<i>corrugicaudatus</i>	[102]
4	<i>Cryoconicus</i>	<i>antiarktos</i>	[109]
5	<i>Dactylobiotus</i>	<i>ovimutans</i>	[134]
6	<i>Diphascion</i>	<i>puchalskii</i>	[133]
7	<i>Diphascion</i>	<i>rudnickii</i>	[133]
8	<i>Hebesuncus</i>	<i>ryani</i>	[99], [106]
9	<i>Hebesuncus</i>	<i>mollispinus</i>	[64]
10	<i>Hypsibius</i>	<i>conwentzii</i>	[133]
11	<i>Mesobiotus</i>	<i>aradasi</i>	[135]
12	<i>Mesobiotus</i>	<i>hilariae</i>	[22]
13	<i>Mesobiotus</i>	<i>krynauwi</i>	[99]
14	<i>Milnesium</i>	<i>validum</i>	[107]
15	<i>Milnesium</i>	<i>rastrum</i>	[136]
16	<i>Milnesium</i>	<i>antarcticum</i>	[100], [103], [137]
17	<i>Milnesium</i>	<i>quadrifidum</i> cv	[64]
18	<i>Mixibius</i>	<i>felix</i>	[107]
19	<i>Mopsechiniscus</i>	<i>franciscae</i>	[108]
20	<i>Paramacrobotus</i>	<i>fairbanksi</i>	[64]
21	<i>Pseudechiniscus</i>	<i>titianae</i>	[22], [106]
22	<i>Ramajendas</i>	<i>dastychi</i>	[138]
23	<i>Ramazzottius</i>	<i>sabatinae</i>	[109]

Some experiments related to short-duration flights demonstrated the ability of tardigrades to survive in an anhydrobiotic state in open space environments in low-Earth orbit [47], [139]. Furthermore, tardigrades offer valuable insights into the origin and evolution of life, enriching our understanding of the intricate tree of life on Earth and shedding light on the potential for life to arise in diverse

environments [47]. The study of tardigrades provides valuable insights into organisms' adaptability and survival strategies. Their unique characteristics make them fascinating objects of study that contribute to expanding our understanding of the potential for life in the universe [140].

#### VII. TARDIGRADES AND THEIR POTENTIAL APPLICATIONS

The study of tardigrades is of great importance for several reasons. First, tardigrades exhibit remarkable survivability, which allows them to withstand extreme conditions such as desiccation, high radiation, and temperature extremes. By delving deeper into the mechanisms driving their exceptional resilience, valuable insights can be gained into adaptations and survival strategies in hostile environments [1]. Tardigrades, such as *Macrobotus hufelandi*, are well known for their remarkable temperature tolerance, showing resistance to temperatures above 100°C for up to 30 min [30]. *Ramazzottius varieornatus* is also classified as a highly tolerant species [141], and *Richtersius coronifer* presents a resistance to high temperature, making both particularly intriguing in their adaptation to global warming [142]. Exploring cryptobiosis and anhydrobiosis mechanisms holds excellent promise for diverse fields, such as medicine, biotechnology, and stress-resistant plant breeding [42].

On the other hand, Antarctica is a relatively unexplored habitat, with limited knowledge about their biodiversity, despite harboring a great variety of unicellular and multicellular organisms. Tardigrades, in particular, play an essential role in these ecosystems, and their study contributes to a better understanding of their ecological dynamics and functions within these niches [143]. For example, limnoterrestrial tardigrades play crucial roles within Antarctic ecosystems, where terrestrial microinvertebrates provide carbon and nutrient cycling roles in soil environments, as a consequence of the absent of larger macroinvertebrates [144]. This role becomes particularly relevant in the absence of other regulatory agents, highlighting the importance of these tardigrades in maintaining ecological balance [145], [146]. The study of these ecological functions not only guides the development of appropriate conservation strategies but also improves our understanding of the factors influencing the dispersal and distribution of organisms in glacial ecosystems [147].

Finally, tardigrades could play a vital role in the field of health in the future. Such is the case of *Ramazzottius varieornatus*, one of the tardigrades mentioned above and one of the most tolerant to stressful environmental conditions. From this organism, it was possible to extract a protein exclusively associated with tardigrades, which, when added to a human cell culture and subjected to X-rays and dehydration, generated a cell culture that was 40% more radioresistant and tolerant to water deficit [62]. The above highlights an existing potential in tardigrade proteins as gene protectors.

## VIII. CONCLUSION

Antarctica is a continent with very extreme conditions [148], [149], [150], [151]; [152], [153]; it is an ideal place for the study of tardigrades. These are extraordinarily resilient organisms. Their ability to survive in extreme conditions such as desiccation, radiation, and extreme temperatures makes these organisms tolerant and resistant to extreme ecosystems such as Antarctica. They are interesting candidates for understanding the possibility of life on other planets making these organisms a coveted model for astrobiology research. Also tardigrades can enter cryptobiosis and anhydrobiosis holds promising prospects in fields such as medicine and biotechnology. Studying tardigrades in Antarctica can provide insights into how microorganisms adapt to and influence geological processes and could be a key factor in understanding climate change and their role in glacial ecosystems. The Antarctic Peninsula has a higher variety of tardigrade species than continental Antarctica [148]. However, knowledge of the biogeographic distribution, abundance, and diversity of tardigrades in Antarctica remains limited making them an attractive research topic. The relationships between tardigrades, geology, and space research have been even less studied, but with a great perspective of development as a research topic. For this reason, the present article aims to review the research potential of limno-terrestrial tardigrades in Antarctica as a model for geobiological and astrobiological studies.

## ACKNOWLEDGMENT

Acknowledgments. The authors thank the Directorate of Research of the Universidad Tecnológica de Bolívar for financing this project, and the Colombian Ocean Commission coordinator of the Colombian Antarctic program.

## REFERENCES

- [1] N. Møbjerg and R. C. Neves, 'New insights into survival strategies of tardigrades', *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, vol. 254, p. 110890, Apr. 2021, doi: 10.1016/j.cbpa.2020.110890.
- [2] K. Wałach and B. Blagden, 'Tardigrade stowaways: Literature review of Propyxidium tardigradum (Ciliophora, Peritrichia) and its first record in Scotland', *European Journal of Protistology*, vol. 89, p. 125974, Jun. 2023, doi: 10.1016/j.ejop.2023.125974.
- [3] M. Jezierska *et al.*, 'Oogenesis in the tardigrade *Hypsibius exemplaris* Gąsiorek, Stec, Morek & Michalczyk, 2018 (Eutardigrada, Hypsibiidae)', *Micron*, vol. 150, p. 103126, Nov. 2021, doi: 10.1016/j.micron.2021.103126.
- [4] N. Guil, 'New records and within-species variability of Iberian tardigrades (Tardigrada), with comments on the species from the *Echiniscus blumi-canadensis* series', *Zootaxa*, vol. 1757, no. 1, p. 1, Apr. 2008, doi: 10.11646/zootaxa.1757.1.1.
- [5] D. R. Nelson, R. Guidetti, and L. Rebecchi, 'Chapter 17 - Phylum Tardigrada', in *Thorpe and Covich's Freshwater Invertebrates (Fourth Edition)*, J. H. Thorpe and D. C. Rogers, Eds., Boston: Academic Press, 2015, pp. 347–380. doi: 10.1016/B978-0-12-385026-3.00017-6.
- [6] K. Arakawa, 'No evidence for extensive horizontal gene transfer from the draft genome of a tardigrade', *Proceedings of the National*

- Academy of Sciences*, vol. 113, no. 22, pp. E3057–E3057, May 2016, doi: 10.1073/pnas.1602711113.
- [7] K. A. S. Schultze, *Macrobiosus Hufelandii, animal e crustaceorum classe novum, reviviscendi post diuturnam asphyxiam et ariditatem potens*. Curths, 1834.
- [8] M. Vecchi, A. Tsvetkova, D. Stec, C. Ferrari, S. Calhim, and D. Tumanov, 'Expanding *Acutuncus*: Phylogenetics and morphological analyses reveal a considerably wider distribution for this tardigrade genus', *Molecular Phylogenetics and Evolution*, vol. 180, p. 107707, Mar. 2023, doi: 10.1016/j.ympev.2023.107707.
- [9] D. R. Nelson, P. J. Bartels, and N. Guil, 'Tardigrade Ecology', in *Water Bears: The Biology of Tardigrades*, R. O. Schill, Ed., in *Zoological Monographs*, Cham: Springer International Publishing, 2018, pp. 163–210. doi: 10.1007/978-3-319-95702-9\_7.
- [10] H. Kumagai, K. Kondo, and T. Kunieda, 'Application of CRISPR/Cas9 system and the preferred no-indel end-joining repair in tardigrades', *Biochemical and Biophysical Research Communications*, vol. 623, pp. 196–201, Oct. 2022, doi: 10.1016/j.bbrc.2022.07.060.
- [11] È. Beltrán-Pardo and J. Bernal-Villegas, 'Acercamiento a los estudios actuales sobre el filo Tardigrada y su importancia en la medicina', *Universitas Médica*, vol. 50, no. 3, pp. 380–403, 2009.
- [12] D. Keilin, 'The Leeuwenhoek Lecture - The problem of anabiosis or latent life: history and current concept', *Proceedings of the Royal Society of London. Series B - Biological Sciences*, vol. 150, no. 939, pp. 149–191, Jan. 1997, doi: 10.1098/rspb.1959.0013.
- [13] R. O. Schill and G. B. Fritz, 'Desiccation tolerance in embryonic stages of the tardigrade', *Journal of Zoology*, vol. 276, no. 1, pp. 103–107, 2008, doi: 10.1111/j.1469-7998.2008.00474.x.
- [14] T. Altiero, I. Giovannini, R. Guidetti, and L. Rebecchi, 'Life history traits and reproductive mode of the tardigrade *Acutuncus antarcticus* under laboratory conditions: strategies to colonize the Antarctic environment', *Hydrobiologia*, vol. 761, no. 1, pp. 277–291, Nov. 2015, doi: 10.1007/s10750-015-2315-0.
- [15] K. Zawierucha, M. Kolicka, N. Takeuchi, and Ł. Kaczmarek, 'What animals can live in cryoconite holes? A faunal review', *Journal of Zoology*, vol. 295, no. 3, pp. 159–169, 2015, doi: 10.1111/jzo.12195.
- [16] L. Jadhav, V. Phalke, S. Panse, S. Patil, and A. Bankar, 'Chapter 12 - Biodiversity of cold-adapted extremophiles from Antarctica and their biotechnological potential', in *Microbial Diversity in Hotspots*, A. Gunjal and S. Shinde, Eds., Academic Press, 2022, pp. 231–265. doi: 10.1016/B978-0-323-90148-2.00013-4.
- [17] F. Richters, 'Vorläufiger Bericht über die antarktische Moosfauna. Verhandlungen der Deutschen Zoologischen Gesellschaft', 1904, pp. 236–239. Accessed: Sep. 22, 2024. [Online]. Available: <https://www.marinespecies.org/tardigrada/aphia.php?p=sourcedetail&id=240648>
- [18] F. Richters, *Wissenschaftliche Ergebnisse der Schwedischen Südpolar-Expedition, 1901-1903*, vol. Bd. V. Liefg 2. Stockholm, 1908. [Online]. Available: <https://www.biodiversitylibrary.org/bibliography/6756>
- [19] J. Murray, 'XII.—Scottish National Antarctic Expedition: Tardigrada of the South Orkneys', *Earth and Environmental Science Transactions of The Royal Society of Edinburgh*, vol. 45, no. 2, pp. 323–334, Jan. 1907, doi: 10.1017/S0080456800022754.
- [20] J. Murray, *British Antarctic Expedition, 1907-9, under the command of Sir E.H. Shackleton, c.v.o. Reports on the scientific investigations*, vol. Vol. 1 Biology (Part V). London, 1910. [Online]. Available: <https://www.biodiversitylibrary.org/bibliography/22427>
- [21] P. Convey and S. J. McInnes, 'Exceptional Tardigrade-Dominated Ecosystems in Ellsworth Land, Antarctica', *Ecology*, vol. 86, no. 2, pp. 519–527, 2005, doi: 10.1890/04-0684.
- [22] M. Vecchi *et al.*, 'Integrative systematic studies on tardigrades from Antarctica identify new genera and new species within Macrotoleida and Echiniscoidea', *Invert. Systematics*, vol. 30, no. 4, pp. 303–322, Aug. 2016, doi: 10.1071/IS15033.
- [23] A. Polishchuk, P. Kayastha, P. Kovalenko, I. Parnikoza, and Ł. Kaczmarek, 'New Records of Tardigrades from the Danco and

- Graham Coasts, the Maritime Antarctic', *annz*, vol. 73, no. 1, pp. 17–28, Mar. 2023, doi: 10.3161/00034541ANZ2023.73.1.002.
- [24] K. A. Short, 'Life in the extreme: when did tardigrades colonise Antarctica?', Ph.D., University of Bristol, 2021. Accessed: Jul. 09, 2023. [Online]. Available: <http://hdl.handle.net/1983/9fdb8089-327e-41be-835c-346282935ef9>
- [25] W. Morek, J. A. Ciosek, and Ł. Michalczyk, 'Description of Milnesium pentapapillatum sp. nov., with an amendment of the diagnosis of the order Apochela and abolition of the class Apotardigrada (Tardigrada)', *Zoologischer Anzeiger*, vol. 288, pp. 107–117, Sep. 2020, doi: 10.1016/j.jcz.2020.07.002.
- [26] M. Czerneková, K. Janelt, S. Student, K. I. Jönsson, and I. Poprawa, 'A comparative ultrastructure study of storage cells in the eutardigrade Richtersius coronifer in the hydrated state and after desiccation and heating stress', *PLOS ONE*, vol. 13, no. 8, p. e0201430, Aug. 2018, doi: 10.1371/journal.pone.0201430.
- [27] V. Gross, S. Treffkorn, J. Reichelt, L. Epple, C. Lüter, and G. Mayer, 'Miniaturization of tardigrades (water bears): Morphological and genomic perspectives', *Arthropod Structure & Development*, vol. 48, pp. 12–19, Jan. 2019, doi: 10.1016/j.asd.2018.11.006.
- [28] M. Tsujimoto, S. Imura, and H. Kanda, 'Recovery and reproduction of an Antarctic tardigrade retrieved from a moss sample frozen for over 30 years', *Cryobiology*, vol. 72, no. 1, pp. 78–81, Feb. 2016, doi: 10.1016/j.cryobiol.2015.12.003.
- [29] J. S. Clegg, 'Cryptobiosis — a peculiar state of biological organization', *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, vol. 128, no. 4, pp. 613–624, Apr. 2001, doi: 10.1016/S1096-4959(01)00300-1.
- [30] R. C. Neves, A. Møbjerg, M. Kodama, J. Ramos-Madrigal, M. T. P. Gilbert, and N. Møbjerg, 'Differential expression profiling of heat stressed tardigrades reveals major shift in the transcriptome', *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, vol. 267, p. 111169, May 2022, doi: 10.1016/j.cbpa.2022.111169.
- [31] J. D. Hibshman, J. S. Clegg, and B. Goldstein, 'Mechanisms of Desiccation Tolerance: Themes and Variations in Brine Shrimp, Roundworms, and Tardigrades', *Frontiers in Physiology*, vol. 11, 2020, Accessed: Jul. 07, 2023. [Online]. Available: <https://www.frontiersin.org/articles/10.3389/fphys.2020.592016>
- [32] K. A. Halberg, A. Jørgensen, and N. Møbjerg, 'Desiccation Tolerance in the Tardigrade Richtersius coronifer Relies on Muscle Mediated Structural Reorganization', *PLOS ONE*, vol. 8, no. 12, p. e85091, Dec. 2013, doi: 10.1371/journal.pone.0085091.
- [33] A. Pigoñ and B. Weglarska, 'Rate of Metabolism in Tardigrades during Active Life and Anabiosis', *Nature*, vol. 176, no. 4472, Art. no. 4472, Jul. 1955, doi: 10.1038/176121b0.
- [34] B. H. Pedersen, H. Malte, H. Ramløv, and K. Finster, 'A method for studying the metabolic activity of individual tardigrades by measuring oxygen uptake using microrespirometry', *Journal of Experimental Biology*, vol. 223, no. 22, p. jeb233072, Nov. 2020, doi: 10.1242/jeb.233072.
- [35] B. H. Pedersen, H. Malte, K. Finster, and H. Ramløv, 'Respiration Measurements of Individual Tardigrades of the Species Richtersius cf. coronifer as a Function of Temperature and Salinity and Termination of Anhydrobiosis', *Astrobiology*, vol. 21, no. 7, pp. 853–865, Jul. 2021, doi: 10.1089/ast.2020.2371.
- [36] E. Beltrán-Pardo, K. I. Jönsson, M. Harms-Ringdahl, S. Haghdoost, and A. Wojcik, 'Tolerance to Gamma Radiation in the Tardigrade Hypsibius dujardini from Embryo to Adult Correlate Inversely with Cellular Proliferation', *PLOS ONE*, vol. 10, no. 7, p. e0133658, Jul. 2015, doi: 10.1371/journal.pone.0133658.
- [37] K. I. Jönsson, S. Borsari, and L. Rebecchi, 'Anhydrobiotic Survival in Populations of the Tardigrades Richtersius coronifer and Ramazzottius oberhaeuseri from Italy and Sweden', *Zoologischer Anzeiger - A Journal of Comparative Zoology*, vol. 240, no. 3, pp. 419–423, Jan. 2001, doi: 10.1078/0044-5231-00050.
- [38] L. Rebecchi, T. Altiero, and R. Guidetti, 'Anhydrobiosis: the extreme limit of desiccation tolerance', *Invertebrate Survival Journal*, vol. 4, no. 2, Art. no. 2, Jun. 2007, Accessed: Feb. 01, 2024. [Online]. Available: <https://www.isj.unimore.it/index.php/ISJ/article/view/144>
- [39] S. Hengherr, M. R. Worland, A. Reuner, F. Brümmer, and R. O. Schill, 'Freeze tolerance, supercooling points and ice formation: comparative studies on the subzero temperature survival of limno-terrestrial tardigrades', *Journal of Experimental Biology*, vol. 212, no. 6, pp. 802–807, Mar. 2009, doi: 10.1242/jeb.025973.
- [40] S. Hengherr, M. R. Worland, A. Reuner, F. Brümmer, and R. O. Schill, 'High-Temperature Tolerance in Anhydrobiotic Tardigrades Is Limited by Glass Transition', *Physiological and Biochemical Zoology*, vol. 82, no. 6, pp. 749–755, Nov. 2009, doi: 10.1086/605954.
- [41] F. Ono *et al.*, 'Effect of ultra-high pressure on small animals, tardigrades and Artemia', *Cogent Physics*, vol. 3, no. 1, p. 1167575, Dec. 2016, doi: 10.1080/23311940.2016.1167575.
- [42] A. Møbjerg *et al.*, 'Extreme freeze-tolerance in cryophilic tardigrades relies on controlled ice formation but does not involve significant change in transcription', *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, vol. 271, p. 111245, Sep. 2022, doi: 10.1016/j.cbpa.2022.111245.
- [43] L. K. B. Hvidepil and N. Møbjerg, 'New insights into osmobiosis and chemobiosis in tardigrades', *Front Physiol*, vol. 14, p. 1274522, Oct. 2023, doi: 10.3389/fphys.2023.1274522.
- [44] T. L. Hygum, L. K. B. Clausen, K. A. Halberg, A. Jørgensen, and N. Møbjerg, 'Tun formation is not a prerequisite for desiccation tolerance in the marine tidal tardigrade Echiniscoides sigmundi', *Zoological Journal of the Linnean Society*, vol. 178, no. 4, pp. 907–911, Dec. 2016, doi: 10.1111/zoj.12444.
- [45] K. I. Jönsson, E. B. Levine, A. Wojcik, S. Haghdoost, and M. Harms-Ringdahl, 'Environmental Adaptations: Radiation Tolerance', in *Water Bears: The Biology of Tardigrades*, R. O. Schill, Ed., in Zoological Monographs. , Cham: Springer International Publishing, 2018, pp. 311–330. doi: 10.1007/978-3-319-95702-9\_12.
- [46] K. I. Jönsson, 'Radiation Tolerance in Tardigrades: Current Knowledge and Potential Applications in Medicine', *Cancers*, vol. 11, no. 9, Art. no. 9, Sep. 2019, doi: 10.3390/cancers11091333.
- [47] K. I. Jönsson, 'Tardigrades as a Potential Model Organism in Space Research', *Astrobiology*, vol. 7, no. 5, pp. 757–766, Oct. 2007, doi: 10.1089/ast.2006.0088.
- [48] K. I. Jönsson, E. Rabbow, R. O. Schill, M. Harms-Ringdahl, and P. Rettberg, 'Tardigrades survive exposure to space in low Earth orbit', *Current Biology*, vol. 18, no. 17, pp. R729–R731, Sep. 2008, doi: 10.1016/j.cub.2008.06.048.
- [49] T. L. Hygum *et al.*, 'Comparative Investigation of Copper Tolerance and Identification of Putative Tolerance Related Genes in Tardigrades', *Frontiers in Physiology*, vol. 8, 2017, Accessed: Feb. 03, 2024. [Online]. Available: <https://www.frontiersin.org/articles/10.3389/fphys.2017.00095>
- [50] O. O. Ojekunle and A. Sodipe, 'Antioxidative Effect of Selenium in Cadmium-Exposed Tardigrade (H. exemplaris)', *Water Air Soil Pollut*, vol. 231, no. 12, p. 577, Dec. 2020, doi: 10.1007/s11270-020-04938-8.
- [51] J. F. Fleming, D. Pisani, and K. Arakawa, 'The Evolution of Temperature and Desiccation-Related Protein Families in Tardigrada Reveals a Complex Acquisition of Extremotolerance', *Genome Biology and Evolution*, vol. 16, no. 1, p. evad217, Jan. 2024, doi: 10.1093/gbe/evad217.
- [52] Ł. Kaczmarek *et al.*, 'How temperature affects the body size of terrestrial tardigrades', *Zoological Journal of the Linnean Society*, vol. 200, no. 1, pp. 144–155, Jan. 2024, doi: 10.1093/zoolinnean/zlad098.
- [53] D. D. Horikawa, K. Kawai, S. Koseki, K. Iwata, T. Okuda, and K. Yamamoto, 'High Hydrostatic Pressure Tolerance of Tardigrades', *高圧バイオサイエンスとバイオテクノロジー*, vol. 1, no. 1, pp. 157–160, 2007, doi: 10.11229/hpbb.1.157.
- [54] N. Kasianchuk, P. Rzymiski, and Ł. Kaczmarek, 'The biomedical potential of tardigrade proteins: A review', *Biomedicine &*

- Pharmacotherapy*, vol. 158, p. 114063, Feb. 2023, doi: 10.1016/j.biopha.2022.114063.
- [55] P. Hagelbäck and K. I. Jönsson, 'An experimental study on tolerance to hypoxia in tardigrades', *Front Physiol*, vol. 14, p. 1249773, Sep. 2023, doi: 10.3389/fphys.2023.1249773.
- [56] L. Rebecchi, M. Cesari, T. Altiero, A. Frigieri, and R. Guidetti, 'Survival and DNA degradation in anhydrobiotic tardigrades', *Journal of Experimental Biology*, vol. 212, no. 24, pp. 4033–4039, Dec. 2009, doi: 10.1242/jeb.033266.
- [57] J. Ingels, D. Zeppilli, and O. Giere, 'Meiofauna—Adapted to Life at the Limits', in *New Horizons in Meiofauna Research: Profiles, Patterns and Potentials*, O. Giere and M. Schratzberger, Eds., Cham: Springer International Publishing, 2023, pp. 363–400. doi: 10.1007/978-3-031-21622-0\_11.
- [58] I. Giovannini, T. C. Boothby, M. Cesari, B. Goldstein, R. Guidetti, and L. Rebecchi, 'Production of reactive oxygen species and involvement of bioprotectants during anhydrobiosis in the tardigrade *Paramacrobiotus spatialis*', *Sci Rep*, vol. 12, no. 1, Art. no. 1, Feb. 2022, doi: 10.1038/s41598-022-05734-6.
- [59] C. Chavez, G. Cruz-Becerra, J. Fei, G. A. Kassavetis, and J. T. Kadonaga, 'The tardigrade damage suppressor protein binds to nucleosomes and protects DNA from hydroxyl radicals', *eLife*, vol. 8, p. e47682, Oct. 2019, doi: 10.7554/eLife.47682.
- [60] R. D. Escarcega *et al.*, 'The Tardigrade damage suppressor protein Dsup promotes DNA damage in neurons', *Molecular and Cellular Neuroscience*, vol. 125, p. 103826, Jun. 2023, doi: 10.1016/j.mcn.2023.103826.
- [61] R. Aguilar *et al.*, 'Multivalent binding of the tardigrade Dsup protein to chromatin promotes yeast survival and longevity upon exposure to oxidative damage', *Res Sq*, p. rs.3.rs-3182883, Jul. 2023, doi: 10.21203/rs.3.rs-3182883/v1.
- [62] T. Hashimoto *et al.*, 'Extremotolerant tardigrade genome and improved radiotolerance of human cultured cells by tardigrade-unique protein', *Nat Commun*, vol. 7, no. 1, Art. no. 1, Sep. 2016, doi: 10.1038/ncomms12808.
- [63] A. Yamaguchi *et al.*, 'Two Novel Heat-Soluble Protein Families Abundantly Expressed in an Anhydrobiotic Tardigrade', *PLOS ONE*, vol. 7, no. 8, p. e44209, Aug. 2012, doi: 10.1371/journal.pone.0044209.
- [64] L. Kaczmarek *et al.*, 'New Records of Antarctic Tardigrada with Comments on Interpopulation Variability of the *Paramacrobiotus fairbanksi* Schill, Förster, Dandekar and Wolf, 2010', *Diversity*, vol. 12, no. 3, Art. no. 3, Mar. 2020, doi: 10.3390/d12030108.
- [65] F. Richters, *Zur Kenntnis der Tardigraden*, 28, 1904.
- [66] K. Morikawa, 'Notes on some Tardigrada from the Antarctic region', *SPECIAL PUBLICATIONS FROM THE SETO MARINE BIOLOGICAL LABORATORY*, vol. 1, no. 17, pp. 1–6, Aug. 1962, Accessed: Oct. 15, 2022. [Online]. Available: <https://repository.kulib.kyoto-u.ac.jp/dspace/handle/2433/176447>
- [67] W. R. Miller, H. Heatwole, R. W. J. Pidgeon, and G. R. Gardiner, 'Tardigrades of the Australian Antarctic Territories: The Larsemann Hills, East Antarctica', *Transactions of the American Microscopical Society*, vol. 113, no. 2, pp. 142–160, 1994, doi: 10.2307/3226642.
- [68] J. D. Miller, P. Hornet, H. Heatwole, W. R. Miller, and L. Bridges, 'A survey of the terrestrial Tardigrada of the Vestfold Hills, Antarctica', *Hydrobiologia*, vol. 165, no. 1, pp. 197–208, Aug. 1988, doi: 10.1007/BF00025588.
- [69] F. Richters, 'Vorläufiger Bericht über die antarktische Moosfauna', *Verhandlungen der Deutschen Zoologischen Gesellschaft*, vol. 1904, pp. 236–239, 1904.
- [70] Y. Shishida and Y. Ohyama, 'A note on the terrestrial nematodes around Syowa Station, Antarctica', 1986.
- [71] C. W. Thomas, 'On Populations in Antarctic Meltwater Pools', Oct. 1965, Accessed: Jul. 16, 2023. [Online]. Available: <http://hdl.handle.net/10125/7375>
- [72] P. Jennings, 'Tardigrada from the Antarctic Peninsula and Scotia Ridge region', *British Antarctic Survey Bulletin*, vol. 44, pp. 77–95, 1976.
- [73] B. Sohlenius, S. Boström, and A. Hirschfelder, 'Distribution patterns of microfauna (nematodes, rotifers and tardigrades) on nunataks in Dronning Maud Land, East Antarctica', *Polar Biol*, vol. 16, no. 3, pp. 191–200, Mar. 1996, doi: 10.1007/BF02329207.
- [74] H. Dastych, 'The Tardigrada from Antarctic with descriptions of several new species', *Acta Zoologica Cracoviensia*, vol. 27, no. 2 (12-21), 1984, Accessed: Oct. 15, 2022. [Online]. Available: <http://agro.icm.edu.pl/agro/element/bwmeta1.element.agro-ac18a5cc-47d5-4de5-a912-c698ed8476ea>
- [75] H. Dastych, 'Redescription of *Hypsibius antarcticus* (Richters, 1904), with some notes on *Hypsibius arcticus* (Murray, 1907)(Tardigrada)', *Mitteilungen aus dem hamburgischen zoologischen Museum und Institut*, vol. 88, pp. 141–159, 1991.
- [76] G. Pilato and M. G. Binda, 'Acutuncus, a new genus of Hypsibiidae (Eutardigrada)', *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg*, vol. 12, no. 155, pp. 159–162, 1997.
- [77] G. Pilato and M. G. Binda, 'Three new species of Diphason of the pingue group (Eutardigrada, Hypsibiidae) from Antarctica', *Polar Biol*, vol. 21, no. 5, pp. 335–342, Apr. 1999, doi: 10.1007/s0030000050370.
- [78] M. G. Binda and G. Pilato, 'Diphason (*Adropion*) *tricuspidatum*, a new species of eutardigrade from Antarctica', *Polar Biol*, vol. 23, no. 1, pp. 75–76, Jan. 2000, doi: 10.1007/s0030000050010.
- [79] W. R. Miller, D. S. Horning, and H. F. Heatwole, 'Tardigrades of the Australian Antarctic: Macquarie Island, sub-Antarctica', *Zoologischer Anzeiger - A Journal of Comparative Zoology*, vol. 240, no. 3, pp. 475–491, Jan. 2001, doi: 10.1078/0044-5231-00057.
- [80] W. Miller, D. Horning, and H. Dastych, *Tardigrades of the Australian Antarctic: description of two new species from Macquarie Island, Subantarctica*. 1995.
- [81] H. Dastych and A. E. Drummond, 'Notes on limnic water-bears (Tardigrada) from the Robertskollen nunataks, Dronning Maud Land, Antarctica', *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg*, vol. 12, no. 154, pp. 111–117, 1996.
- [82] K. Utsugi and Y. Ohyama, 'Antarctic Tardigrada II. *Molodezhnaya* and Mt. Riiser-Larsen areas', presented at the Proceedings of the NIPR Symposium on Polar Biology, 1991, pp. 161–170.
- [83] L. Sømme and T. Meier, 'Cold tolerance in Tardigrada from Dronning Maud Land, Antarctica', *Polar Biol*, vol. 15, no. 3, pp. 221–224, Feb. 1995, doi: 10.1007/BF00239062.
- [84] K. Utsugi and Y. Ohyama, 'Antarctic tardigrada 111. Fildes peninsula of king george island', presented at the Proc. NIPR Symp. Polar Biol, Citeseer, 1993, pp. 139–151.
- [85] S. J. McInnes, 'Tardigrades from Signy Island, South Orkney Islands, with particular reference to freshwater species', *Journal of Natural History*, vol. 29, no. 6, pp. 1419–1445, Dec. 1995, doi: 10.1080/00222939500770601.
- [86] P. Jennings, 'The Tardigrada of Signy Island, South Orkney Islands, with a note on the Rotifera', *British Antarctic Survey Bulletin*, vol. 44, pp. 1–25, 1976.
- [87] Utsugi K. and Ohyama Y., 'Antarctic Tardigrada', vol. 2, pp. 190–197, Sep. 1989, Accessed: Dec. 06, 2022. [Online]. Available: [https://nipr.repo.nii.ac.jp/index.php?active\\_action=repository\\_view\\_main\\_item\\_detail&page\\_id=13&block\\_id=104&item\\_id=5066&item\\_no=1](https://nipr.repo.nii.ac.jp/index.php?active_action=repository_view_main_item_detail&page_id=13&block_id=104&item_id=5066&item_no=1)
- [88] H. Dastych, P. Ryan, and B. Watkins, 'Notes on Tardigrada from western Dronning Maud Land (Antarctica) with a description of two new species', *Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg*, vol. 10, no. 139/140, pp. 57–66, 1990.
- [89] M. Sudzuki, 'On the microfauna of the Antarctic region III. Microbiota of the terrestrial interstrices', 1979.
- [90] M. Sudzuki, 'On the Microfauna of the Antarctic Region: I. Moss-water community at Langhovde', 1964.
- [91] J. Murray, 'XXVII.—The Tardigrada of the Scottish Lochs', *Earth and Environmental Science Transactions of The Royal Society of Edinburgh*, vol. 41, no. 3, pp. 677–698, 1906.
- [92] W. R. Miller and H. Heatwole, 'Tardigrades of the Australian Antarctic Territories: The Mawson Coast, East Antarctica', *Invertebrate Biology*, vol. 114, no. 1, pp. 27–38, 1995, doi: 10.2307/3226950.

- [93] W. R. Miller, H. Heatwole, R. W. J. Pidgeon, and G. R. Gardiner, 'Tardigrades of the Australian Antarctic Territories: The Larsemann Hills, East Antarctica', *Transactions of the American Microscopical Society*, vol. 113, no. 2, pp. 142–160, 1994, doi: 10.2307/3226642.
- [94] W. Miller, J. Miller, and H. Heatwole, 'Tardigrades of the Australian Antarctic Territories: The Windmill Islands, East Antarctica', *Zoological Journal of the Linnean Society*, vol. 116, no. 1–2, pp. 175–184, 1996.
- [95] G. Ramazzotti, 'Tardigradi delle Isole Kerguelen e descrizione della nuova specie *Hypsibius* (L.) renaudi', *Mem Ist Ital Idrobiol*, vol. 29, pp. 141–144, 1972.
- [96] E. C. Dougherty and L. G. Harris, 'Antarctic Micrometazoa: Fresh-Water Species in the McMurdo Sound Area', *Science*, vol. 140, no. 3566, pp. 497–498, May 1963, doi: 10.1126/science.140.3566.497.
- [97] R. B. Heywood, V. E. Fuchs, and R. M. Laws, 'A limnological survey of the Ablation Point area, Alexander Island, Antarctica', *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, vol. 279, no. 963, pp. 39–54, Jan. 1997, doi: 10.1098/rstb.1977.0070.
- [98] B. J. Adams *et al.*, 'Diversity and distribution of Victoria Land biota', *Soil Biology and Biochemistry*, vol. 38, no. 10, pp. 3003–3018, Oct. 2006, doi: 10.1016/j.soilbio.2006.04.030.
- [99] B. Sohlenius, S. Boström, and K. Ingemar Jönsson, 'Occurrence of nematodes, tardigrades and rotifers on ice-free areas in East Antarctica', *Pedobiologia*, vol. 48, no. 4, pp. 395–408, Aug. 2004, doi: 10.1016/j.pedobi.2004.06.001.
- [100] J. Smykla, N. Iakovenko, M. Devetter, and Ł. Kaczmarek, 'Diversity and distribution of tardigrades in soils of Edmonson Point (Northern Victoria Land, continental Antarctica)', *Czech Polar Reports*, vol. 2, no. 2, Art. no. 2, Jun. 2012, doi: 10.5817/CPR2012-2-6.
- [101] J. A. Gibson, L. Cromer, J. T. Agius, S. J. McInnes, and N. J. Marley, 'Tardigrade eggs and exuviae in Antarctic lake sediments: insights into Holocene dynamics and origins of the fauna', *Journal of Limnology*, vol. 66, no. Supple, pp. 65–71, 2007.
- [102] S. J. McInnes, 'Echiniscuscorrugicaudatus (Heterotardigrada; Echiniscidae) a new species from Ellsworth Land, Antarctica', *Polar Biol*, vol. 33, no. 1, pp. 59–70, Jan. 2010, doi: 10.1007/s00300-009-0684-4.
- [103] D. V. Tumanov, 'Five new species of the genus *Milnesium* (Tardigrada, Eutardigrada, Milnesiidae)', *Zootaxa*, vol. 1122, no. 1, Art. no. 1, Feb. 2006, doi: 10.11646/zootaxa.1122.1.1.
- [104] A. Velasco-Castrillón *et al.*, 'Distribution and Diversity of Soil Microfauna from East Antarctica: Assessing the Link between Biotic and Abiotic Factors', *PLOS ONE*, vol. 9, no. 1, p. e87529, Jan. 2014, doi: 10.1371/journal.pone.0087529.
- [105] H. Kagoshima, S. Imura, and A. C. Suzuki, 'Molecular and morphological analysis of an Antarctic tardigrade, *Acutuncus antarcticus*', *Journal of Limnology*, vol. 72, no. s1, Art. no. s1, Mar. 2013, doi: 10.4081/jlimnol.2013.s1.e3.
- [106] M. Tsujimoto, S. J. McInnes, P. Convey, and S. Imura, 'Preliminary description of tardigrade species diversity and distribution pattern around coastal Syowa Station and inland Sør Rondane Mountains, Dronning Maud Land, East Antarctica', *Polar Biol*, vol. 37, no. 9, pp. 1361–1367, Sep. 2014, doi: 10.1007/s00300-014-1516-8.
- [107] G. Pilato, G. Sabella, V. D'urso, and O. Lisi, 'Two new species of Eutardigrada from Victoria Land, Antarctica', *Zootaxa*, vol. 4317, no. 3, Art. no. 3, Sep. 2017, doi: 10.11646/zootaxa.4317.3.6.
- [108] R. Guidetti, L. Rebecchi, M. Cesari, and S. J. McInnes, 'Mopsechiniscus franciscae, a new species of a rare genus of Tardigrada from continental Antarctica', *Polar Biol*, vol. 37, no. 9, pp. 1221–1233, Sep. 2014, doi: 10.1007/s00300-014-1514-x.
- [109] R. Guidetti, E. Massa, R. Bertolani, L. Rebecchi, and M. Cesari, 'Increasing knowledge of Antarctic biodiversity: new endemic taxa of tardigrades (Eutardigrada; Ramazzottiidae) and their evolutionary relationships', *Systematics and Biodiversity*, vol. 17, no. 6, pp. 573–593, Aug. 2019, doi: 10.1080/14772000.2019.1649737.
- [110] A. Velasco-Castrillón *et al.*, 'Mitochondrial DNA analyses reveal widespread tardigrade diversity in Antarctica', *Invert. Systematics*, vol. 29, no. 6, pp. 578–590, Dec. 2015, doi: 10.1071/IS14019.
- [111] F. Pereira, J. Carneiro, and A. Amorim, 'Identification of Species with DNA-Based Technology: Current Progress and Challenges', *Recent Patents on DNA & Gene Sequences*, vol. 2, no. 3, pp. 187–200, Nov. 2008, doi: 10.2174/187221508786241738.
- [112] S. Antil *et al.*, 'DNA barcoding, an effective tool for species identification: a review', *Mol Biol Rep*, vol. 50, no. 1, pp. 761–775, Jan. 2023, doi: 10.1007/s11033-022-08015-7.
- [113] P. Gąsiorek, D. Stec, W. Morek, and Ł. Michalczyk, 'An integrative redescription of *Hypsibius dujardini* (Doyère, 1840), the nominal taxon for Hypsibiodea (Tardigrada: Eutardigrada)', *Zootaxa*, vol. 4415, no. 1, Art. no. 1, Apr. 2018, doi: 10.11646/zootaxa.4415.1.2.
- [114] H. Dastych, 'Redescription and revalidation of the sub-Antarctic tardigrade *Hypsibius murrayi* (Richters, 1907) based on the rediscovered type material (Tardigrada, Panarthropoda)', *Entomologie heute*, vol. 30, pp. 95–115, 2018.
- [115] D. V. Tumanov, 'End of a mystery: Integrative approach reveals the phylogenetic position of an enigmatic Antarctic tardigrade genus *Ramajendas* (Tardigrada, Eutardigrada)', *Zoologica Scripta*, vol. 51, no. 2, pp. 217–231, 2022, doi: 10.1111/zsc.12521.
- [116] G. Visconti, 'Geobiology', in *Climate, Planetary and Evolutionary Sciences: A Machine-Generated Literature Overview*, G. Visconti, Ed., Cham: Springer International Publishing, 2021, pp. 317–329, doi: 10.1007/978-3-030-74713-8\_9.
- [117] Y. Dilek, H. Furnes, and K. Muehlenbachs, *Links Between Geological Processes, Microbial Activities & Evolution of Life: Microbes and Geology*. Springer Science & Business Media, 2008.
- [118] P. G. Falkowski, T. Fenchel, and E. F. Delong, 'The Microbial Engines That Drive Earth's Biogeochemical Cycles', *Science*, vol. 320, no. 5879, pp. 1034–1039, May 2008, doi: 10.1126/science.1153213.
- [119] A. Summers Engel, M. L. Porter, B. K. Kinkle, and T. C. Kane, 'Ecological Assessment and Geological Significance of Microbial Communities from Cesspool Cave, Virginia', *Geomicrobiology Journal*, vol. 18, no. 3, pp. 259–274, Jul. 2001, doi: 10.1080/01490450152467787.
- [120] N. Zhu, J. Wang, Y. Wang, S. Li, and J. Chen, 'Differences in geological conditions have reshaped the structure and diversity of microbial communities in oily soils', *Environmental Pollution*, vol. 306, p. 119404, Aug. 2022, doi: 10.1016/j.envpol.2022.119404.
- [121] L. K. Baumgartner *et al.*, 'Sulfate reducing bacteria in microbial mats: Changing paradigms, new discoveries', *Sedimentary Geology*, vol. 185, no. 3, pp. 131–145, Mar. 2006, doi: 10.1016/j.sedgeo.2005.12.008.
- [122] B. B. Jørgensen and A. Boetius, 'Feast and famine — microbial life in the deep-sea bed', *Nat Rev Microbiol*, vol. 5, no. 10, Art. no. 10, Oct. 2007, doi: 10.1038/nrmicro1745.
- [123] B. Zhang *et al.*, 'Microbial Community Responses to Vanadium Distributions in Mining Geological Environments and Bioremediation Assessment', *Journal of Geophysical Research: Biogeosciences*, vol. 124, no. 3, pp. 601–615, 2019, doi: 10.1029/2018JG004670.
- [124] V. S. Smetacek, 'Role of sinking in diatom life-history cycles: ecological, evolutionary and geological significance', *Mar. Biol.*, vol. 84, no. 3, pp. 239–251, Jan. 1985, doi: 10.1007/BF00392493.
- [125] J. E. Barrett *et al.*, 'Co-variation in soil biodiversity and biogeochemistry in northern and southern Victoria Land, Antarctica', *Antarctic Science*, vol. 18, no. 4, pp. 535–548, Dec. 2006, doi: 10.1017/S0954102006000587.
- [126] D. L. Moorhead, J. E. Barrett, R. A. Virginia, D. H. Wall, and D. Porazinska, 'Organic matter and soil biota of upland wetlands in Taylor Valley, Antarctica', *Polar Biol*, vol. 26, no. 9, pp. 567–576, Sep. 2003, doi: 10.1007/s00300-003-0524-x.
- [127] S. J. McInnes, 'Taxonomy, biodiversity and biogeography: Tardigrada and Antarctic meiofauna', Anglia Ruskin University, 2010. Accessed: Sep. 23, 2024. [Online]. Available: [https://aru.figshare.com/articles/thesis/Taxonomy\\_biodiversity\\_and](https://aru.figshare.com/articles/thesis/Taxonomy_biodiversity_and)

- biogeography\_Tardigarda\_and\_Antarctic\_meiofauna/23752428?file=42192381
- [128] S. J. McInnes and J. Philip, 'An attempt to revisit the global biogeography of limno-terrestrial Tardigrada', *Journal of Limnology*, vol. 66, p. 90, 2007.
- [129] G. Pilato and M. G. Binda, 'Biogeography and Limno-terrestrial Tardigrades: Are They Truly Incompatible Binomials?', *Zoologischer Anzeiger - A Journal of Comparative Zoology*, vol. 240, no. 3, pp. 511–516, Jan. 2001, doi: 10.1078/0044-5231-00061.
- [130] H. Heatwole *et al.*, 'From bedrock to biota: weathering, physico-chemical properties, protozoans and micrometazoans of some soils of East Antarctica', 1999.
- [131] O. O. Ojekunle, 'Morphological Effects of Copper and Cadmium Exposure on Tardigrade (*H. exemplaris*) and the Antioxidative Effect of Selenium', 2020.
- [132] E. Weronika and K. Łukasz, 'Tardigrades in Space Research - Past and Future', *Orig Life Evol Biosph*, vol. 47, no. 4, pp. 545–553, Dec. 2017, doi: 10.1007/s11084-016-9522-1.
- [133] Ł. Kaczmarek *et al.*, 'Tardigrades from *Larus dominicanus* Lichtenstein, 1823 nests on the Argentine Islands (maritime Antarctic)', *Polar Biol*, vol. 41, no. 2, pp. 283–301, Feb. 2018, doi: 10.1007/s00300-017-2190-4.
- [134] J.-H. Kihm *et al.*, 'Integrative description of a new Dactylobiotus (Eutardigrada: Parachela) from Antarctica that reveals an intraspecific variation in tardigrade egg morphology', *Sci Rep*, vol. 10, no. 1, Art. no. 1, Jun. 2020, doi: 10.1038/s41598-020-65573-1.
- [135] M. G. Binda, G. Pilato, and O. Lisi, 'Remarks on Macrobiotus furciger Murray, 1906 and description of three new species of the furciger group (Eutardigrada, Macrobiotidae)', *Zootaxa*, vol. 1075, no. 1, Art. no. 1, Nov. 2005, doi: 10.11646/zootaxa.1075.1.3.
- [136] A. C. Suzuki *et al.*, 'A New Species of Bisexual Milnesium (Eutardigrada: Apochela) Having Aberrant Claws from Innhovde, Dronning Maud Land, East Antarctica', *Jzoo*, vol. 40, no. 3, pp. 246–261, Feb. 2023, doi: 10.2108/zs220085.
- [137] J. Smykla *et al.*, 'Geochemical and biotic factors influencing the diversity and distribution of soil microfauna across ice-free coastal habitats in Victoria Land, Antarctica', *Soil Biology and Biochemistry*, vol. 116, pp. 265–276, Jan. 2018, doi: 10.1016/j.soilbio.2017.10.028.
- [138] Ł. Kaczmarek, K. Janko, J. Smykla, and Ł. Michalczuk, 'Soil tardigrades from the Antarctic Peninsula with a description of a new species and some remarks on the genus Ramajendas (Eutardigrada: Isohypsibiidae)', *Polar Record*, vol. 50, no. 2, pp. 176–182, Apr. 2014, doi: 10.1017/S0032247413000168.
- [139] D. Persson, K. A. Halberg, A. Jørgensen, C. Ricci, N. Møbjerg, and R. M. Kristensen, 'Extreme stress tolerance in tardigrades: surviving space conditions in low earth orbit', *Journal of Zoological Systematics and Evolutionary Research*, vol. 49, no. s1, pp. 90–97, 2011, doi: 10.1111/j.1439-0469.2010.00605.x.
- [140] R. Guidetti, A. M. Rizzo, T. Altiero, and L. Rebecchi, 'What can we learn from the toughest animals of the Earth? Water bears (tardigrades) as multicellular model organisms in order to perform scientific preparations for lunar exploration', *Planetary and Space Science*, vol. 74, no. 1, pp. 97–102, Dec. 2012, doi: 10.1016/j.pss.2012.05.021.
- [141] R. C. Neves, L. K. B. Hvidepil, T. L. Sørensen-Hygun, R. M. Stuart, and N. Møbjerg, 'Thermotolerance experiments on active and desiccated states of *Ramazzottius varieornatus* emphasize that tardigrades are sensitive to high temperatures', *Sci Rep*, vol. 10, no. 1, Art. no. 1, Jan. 2020, doi: 10.1038/s41598-019-56965-z.
- [142] H. Ramløv and P. Westh, 'Cryptobiosis in the Eutardigrade *Adorybiotus (Richtersius) coronifer*: Tolerance to Alcohols, Temperature and *de novo* Protein Synthesis', *Zoologischer Anzeiger - A Journal of Comparative Zoology*, vol. 240, no. 3, pp. 517–523, Jan. 2001, doi: 10.1078/0044-5231-00062.
- [143] K. Zawierucha, J. Buda, T. N. Jaromerska, K. Janko, and P. Gąsiorek, 'Integrative approach reveals new species of water bears (*Pilatobius*, *Grevenius*, and *Acutuncus*) from Arctic cryoconite holes, with the discovery of hidden lineages of *Hypsibius*', *Zoologischer Anzeiger*, vol. 289, pp. 141–165, Nov. 2020, doi: 10.1016/j.jcz.2020.09.004.
- [144] J. S. McCarthy *et al.*, 'Population growth of two limno-terrestrial Antarctic microinvertebrates in different aqueous soil media', *Environ Sci Pollut Res*, vol. 31, no. 22, pp. 33086–33097, May 2024, doi: 10.1007/s11356-024-32905-x.
- [145] S. Sánchez-Moreno, H. Ferris, and N. Guil, 'Role of tardigrades in the suppressive service of a soil food web', *Agriculture, Ecosystems & Environment*, vol. 124, no. 3, pp. 187–192, Apr. 2008, doi: 10.1016/j.agee.2007.09.011.
- [146] M. Tůmová, D. Stec, Ł. Michalczuk, and M. Devetter, 'Buccal tube dimensions and prey preferences in predatory tardigrades', *Applied Soil Ecology*, vol. 170, p. 104303, Feb. 2022, doi: 10.1016/j.apsoil.2021.104303.
- [147] K. Zawierucha, D. Stec, P. K. Dearden, and D. H. Shain, 'Two new tardigrade genera from New Zealand's Southern Alp glaciers display morphological stasis and parallel evolution', *Molecular Phylogenetics and Evolution*, vol. 178, p. 107634, Jan. 2023, doi: 10.1016/j.ympev.2022.107634.
- [148] R. Acevedo-Barrios, Rubiano-Labrador, Carolina, Altamar Mercado, Hernando; Luis Villalba, Jorgea; Monroy-Licht, Andrea; Carranza-López, Liliana; Angélica Leal, María Tovar, David. Identification of tardigrades from the Half Moon Island, Antarctic Proceedings. Proceedings of the LACCEI international Multi-conference for Engineering, Education and Technology2024 22nd LACCEI International Multi-Conference for Engineering, Education and Technology, LACCEI 2024Hybrid, San Jose17 July 2024through 19 July 2024Code 202044of the LACCEI international Multi-conference for Engineering, Education and Technology2024 22nd LACCEI International Multi-Conference for Engineering, Education and Technology, LACCEI 2024Hybrid, San Jose17 July 2024through 19 July 2024Code 202044. DOI 10.18687/LACCEI2024.1.1.445
- [149] R. Acevedo-Barrios, Puentes Martínez, D.A., Hernández Rocha, I.O. et al. Perchlorate in Antarctica, origin, effects, treatments, environmental fate, and astrobiological perspectives: a review. Int. J. Environ. Sci. Technol. (2025). <https://doi.org/10.1007/s13762-024-06004-w>
- [150] R. Acevedo-Barrios, I. Hernández Rocha, D. Puentes Martinez, Rubiano-Labrador C, J. Pasqualino, E. Chavarro-Mesa, A. De La Parra Querra. Psychrobacter sp: perchlorate reducing bacteria, isolated from marine sediments from Margarita Bay, Antarctica. Rosa Leonor. Proceedings of the LACCEI international Multi-conference for Engineering, Education and Technology, 4(13), 16. July-2023. doi:10.18687/LACCEI2023.1.1.995.
- [151] R. Acevedo-Barrios, C. Rubiano-Labrador, D. Navarro-Narvaez, et al. Perchlorate-reducing bacteria from Antarctic marine sediments. Environ Monit Assess 194, 654 2022. doi: 10.1007/s10661-022-10328-w.
- [152] R. Acevedo-Barrios, C. Rubiano-Labrador, y W. Miranda-Castro, "Presence of perchlorate in marine sediments from Antarctica during 2017-2020", Environ. Monit. Assess., vol. 194, núm. 2, p. 102, 2022. <https://doi.org/10.1007/s10661-022-09765-4>
- [153] R. Acevedo-Barrios y J. Olivero-Verbel, "Perchlorate contamination: Sources, effects, and technologies for remediation", Rev. Environ. Contam. Toxicol., vol. 256, pp. 103–120, 2021. [https://doi.org/10.1007/398\\_2021\\_66](https://doi.org/10.1007/398_2021_66)